

Pion and Kaon Condensation at Finite Temperature and Density

Jens O. Andersen*

Department of Physics, Norwegian Institute of Science and Technology, N-7491 Trondheim, Norway

(Dated: February 2, 2008)

In this paper, we study $O(2N)$ -symmetric ϕ^4 -theory at finite temperature and density using the 2PI- $1/N$ expansion. As specific examples, we consider pion condensation at finite isospin chemical potential and kaon condensation at finite chemical potential for hyper charge and isospin charge. We calculate the phase diagrams and the quasiparticle masses for pions and kaons in the large- N limit. It is shown that the effective potential and the gap equation can be renormalized by using local counterterms for the coupling constant and mass parameter, which are independent of temperature and chemical potentials.

PACS numbers: 11.15Bt, 04.25.Nx, 11.10Wx, 12.38Mh

I. INTRODUCTION

$O(N)$ -symmetric field theories are playing a significant role as effective quantum field theories in condensed matter and high-energy physics. For example, the non-relativistic $O(2)$ -symmetric model is used to describe cold atomic Bose gases in the context of Bose-Einstein condensation in harmonic traps [1] and the homogeneous dilute Bose gas [2]. Similarly, the $O(4)$ (non)linear sigma model is used as a low-energy effective theory for QCD. In massless two-flavor QCD, the global symmetry group is $SU(2)_L \times SU(2)_R$ and one then takes advantage of the isomorphism between this group and $O(4)$. This effective theory describes the three pions and the sigma particle. Similarly, the $SU(2)_L \times SU(2)_R$ -symmetric theory is a toy model for the description of kaon condensates [3, 4] in the color-flavor locked phase [5] of high-density QCD.

The thermodynamic functions for hot field theories can be calculated as a series in the coupling constant by resumming certain classes of diagrams from all orders of perturbation theory. However, it turns out that the expansion of these functions in powers of the coupling constant breaks down at temperature unless the coupling is tiny [6, 7, 8]. In order to obtain more stable results, one has developed and applied more advanced resummation techniques, such as screened perturbation theory [9], hard-thermal-loop resummation [10] and resummations based on the two-particle irreducible (2PI) effective action [11, 12, 13, 14]. These approaches all improve the converge of the perturbative series of the thermodynamic quantities.

At large chemical potential and zero temperature, resummation of classes of diagrams is also required [15, 16] to obtain the thermodynamic functions as a series ex-

pansion in the coupling. However, the perturbative series may be better behaved at zero temperature than at high temperature as the plasmon effects are different [17]. In this case, and also at nonzero but low temperature ($T \leq \mu$), hard-dense-loop resummation has proven to be an effective way of reorganizing the calculations [18]. Very recently, Ipp *et al* [19] have presented a new method for calculating the pressure in deconfined QCD at all values of the temperature and the quark chemical potential, which reduces to the known perturbative results in the appropriate limits.

The $1/N$ expansion in conjunction with e.g. the 1PI or 2PI effective-action formalism is another nonperturbative resummation scheme which has been applied at finite temperature [20, 21, 22, 23]. A leading-order calculation typically gives rise to a gap equation for a mass parameter, which is straightforward to solve. At next-to-leading order (NLO), the resulting equations become harder to solve. Due to nontrivial renormalization issues, the NLO problem using the 1PI $1/N$ -expansion, has been solved rigorously only very recently [22]. In particular, it was shown that the thermodynamic potential can be renormalized in a temperature-independent way only at the minimum. In the case of the 2PI formalism there are additional complications since the gap equation for the propagator becomes nonlocal and nontrivial to solve. However, the 2PI formalism is superior when it comes to the description of nonequilibrium phenomena [21].

Renormalization of expressions obtained in resummation schemes such as the 2PI effective-action formalism is a nontrivial issue. In these approximations, one generally resums only certain classes of diagrams from all orders of perturbation theory. Since only a subset of diagrams from a given order in perturbation theory is taken into account, standard proofs of perturbative renormalizability do not apply and one is not guaranteed that ultraviolet divergences can be systematically eliminated. Moreover, it is not obvious that these divergences

*Electronic address: andersen@tf.phys.no

are independent of temperature. Significant progress on these issues has been made in recent years starting with the papers by van Hees and Knoll [24]. They showed that the equation for the two-point function in scalar ϕ^4 -theory can be renormalized by introducing a finite number of counterterms. A more systematic study of the renormalization issues in scalar ϕ^4 -theory was carried out by Blaizot, Iancu and Reinosa [25]. In particular, they focused on the elimination of temperature-dependent divergences. This is sufficient to calculate physical quantities such as the thermodynamic pressure, but only recently has it been shown by Berges *et al* [26] how to fix all the counterterms needed to calculate the proper vertices which are encoded in the effective action. These counterterms are local and they are independent of temperature and chemical potential.

The idea of a Bose-Einstein condensate of pions or kaons in the core of compact stars has a long history and the interest in such condensates is due to their far-reaching consequences for these objects. For example, the presence of a condensate results in enhanced neutrino cooling [27]. Moreover, in contrast to hadronic matter in heavy-ion collisions, bulk matter in compact stars must (on average) be electrically neutral and so a neutrality constraint must be imposed.

In the present paper, we discuss pion and kaon condensation at finite temperature and density using the 2PI-1/ N expansion. Pion condensation and the phase diagram of two-flavor QCD have been investigated using chiral perturbation theory [28, 29, 30, 31], lattice QCD [32, 33], ladder QCD [34], chiral quark model [35], and Nambu Jona-Lasinio models in the mean-field approximation [36, 37, 38]. In Ref. [38], the effects of imposing charge neutrality were studied at zero temperature. There have also been some applications of the linear sigma model at finite temperature and density in either the large- N approximation [37] or the Hartree approximation [39]. In both cases, the quasiparticle masses were calculated, but renormalization issues were not discussed. In the present paper, we generalize part of the finite-temperature calculations of Lenaghan and Rischke [23] by including chemical potentials. More specifically, we study pion and kaon condensation at finite temperature and density using the 2PI 1/ N -expansion paying attention to renormalization issues.

The paper is organized as follows. In Sec. II, we briefly discuss interacting Bose gases at finite density. In Sec. III, we consider pions at finite temperature and isospin chemical potential. We calculate the phase diagram and the quasiparticle masses. In Sec. IV, we investigate kaon condensation in the presence of chemical potentials for isospin and hypercharge. In Sec. V, we summarize and conclude.

II. INTERACTING BOSE GAS

The Euclidean Lagrangian for a Bose gas with N species of massive charged scalars is

$$\mathcal{L} = (\partial_\mu \Phi_i^\dagger)(\partial_\mu \Phi_i) - \frac{H}{\sqrt{2}} (\Phi_1 + \Phi_1^\dagger) + m^2 \Phi_i^\dagger \Phi_i + \frac{\lambda}{2N} (\Phi_i^\dagger \Phi_i)^2, \quad (1)$$

where $i = 1, 2, \dots, N$ and $\Phi_i = (\phi_{2i-1} + i\phi_{2i})/\sqrt{2}$ is a complex field. If $H = 0$, the theory described by Eq. (1) has $(2N - 1)N$ conserved charges which equals the number of generators of the group $O(2N)$. If H is nonzero, the $O(2N)$ symmetry is explicitly broken down to $O(2N - 1)$. A gas of N species of bosons can be characterized by the expectation values of the different conserved charges in addition to the temperature. For each conserved charge Q_i , one may introduce a nonzero chemical potential μ_i . However, it is possible to specify the expectation values of different charges only if they commute. The maximum number of commuting charges is N [40] and these can be chosen as

$$Q_i = \int d^3x j_i^0, \quad (2)$$

where the the current densities j_i^μ are

$$j_i^\mu = \phi_{2i} \partial^\mu \phi_{2i-1} - \phi_{2i-1} \partial^\mu \phi_{2i} \quad (3)$$

The incorporation of a conserved charge Q_i is done by making the substitution

$$\partial_0 \Phi_i \rightarrow (\partial_0 - \mu_i) \Phi_i \quad (4)$$

$$\partial_0 \Phi_i^\dagger \rightarrow (\partial_0 + \mu_i) \Phi_i^\dagger. \quad (5)$$

in the Lagrangian (1). From the path-integral representation of the free energy density $\mathcal{F}$

$$e^{-\beta V \mathcal{F}} = \int \mathcal{D}\Phi_i^* \mathcal{D}\Phi_i e^{-\int_0^\beta d\tau \int d^3x \mathcal{L}}, \quad (6)$$

the expression for the charge density can be written as

$$Q_i = -\frac{\partial \mathcal{F}}{\partial \mu_i}. \quad (7)$$

If we introduce k chemical potentials, the full symmetry group is broken down to $[O(2)]^k \times O(2N - 2k)$. If $m^2 < 0$, the $O(2N)$ symmetry is spontaneously broken down to $O(2N - 1)$. Even if $m^2 > 0$, the symmetry may be broken if one of the chemical potentials, μ_i , is larger than a critical chemical potential $\mu_c = m$. In that case, the $\langle 0 | \phi_{2i-1} | 0 \rangle \neq 0$ and the corresponding $O(2)$ symmetry is broken.

III. PIONS

In this section, we discuss pions at finite chemical potential μ_I for (the third component of the) isospin charge, which corresponds to the current density $j^\mu \sim \phi_2 \partial^\mu \phi_3 - \phi_3 \partial^\mu \phi_2$, and $N = 2$ in Eq. (1). In the vacuum, the $O(4)$ symmetry is broken down to $O(3)$ by a nonzero vacuum expectation value ϕ_0 for ϕ_1 . The symmetry is broken either spontaneously in the chiral limit or explicitly if $H \neq 0$.

We first introduce a nonzero expectation value ρ_0 for ϕ_2 to allow for a charged pion condensate. The field Φ_1 is then written as

$$\Phi_1 = \frac{1}{\sqrt{2}}(\phi_0 + i\rho_0 + \phi_1 + i\phi_2), \quad (8)$$

where ϕ_1 and ϕ_2 are quantum fluctuating fields. The inverse tree-level propagator $D_0^{-1}(\omega_n, p)$ reads

$$D_0^{-1}(\omega_n, p) = \begin{pmatrix} \omega_n^2 + p^2 + m_1^2 & \frac{\lambda}{N}\phi_0\rho_0 & 0 & 0 & \dots \\ \frac{\lambda}{N}\phi_0\rho_0 & \omega_n^2 + p^2 + m_2^2 & -2\mu_I\omega_n & 0 & \dots \\ 0 & 2\mu_I\omega_n & \omega_n^2 + p^2 + m_3^2 & 0 & \dots \\ 0 & 0 & 0 & \omega_n^2 + p^2 + m_4^2 & \dots \\ 0 & 0 & 0 & 0 & \omega_n^2 + p^2 + m_4^2 \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}, \quad (9)$$

where $p = |\mathbf{p}|$, $\omega_n = 2n\pi T$ is the n th Matsubara frequency, and the tree-level masses are

$$m_1^2 = m^2 + \frac{3\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2, \quad (10)$$

$$m_2^2 = -\mu_I^2 + m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{3\lambda}{2N}\rho_0^2, \quad (11)$$

$$m_3^2 = -\mu_I^2 + m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2, \quad (12)$$

$$m_4^2 = m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2. \quad (13)$$

Note that the mass parameter m^2 is negative in the remainder of this section, as we want to break chiral symmetry spontaneously in the chiral limit in the absence of a chemical potential.

The 2PI effective potential can be written as

$$\begin{aligned} \Gamma[\phi_0, \rho_0, D] &= \frac{1}{2}m^2(\phi_0^2 + \rho_0^2) + \frac{\lambda}{8N}(\phi_0^2 + \rho_0^2)^2 \\ &\quad - \frac{1}{2}\mu_I^2\rho_0^2 - H\phi_0 + \frac{1}{2}\text{Tr} \ln D^{-1} \\ &\quad + \frac{1}{2}\text{Tr} D_0^{-1}D + \Phi[D], \end{aligned} \quad (14)$$

where D is the exact propagator and $\Phi[D]$ is the sum of all two-particle irreducible vacuum diagrams. The trace is over space-time as well as field indices. The expectation values ϕ_0 and ρ_0 satisfy the stationarity conditions

$$\frac{\delta\Gamma[\phi_0, \rho_0, D]}{\delta\phi_0} = 0, \quad (15)$$

$$\frac{\delta\Gamma[\phi_0, \rho_0, D]}{\delta\rho_0} = 0. \quad (16)$$

The exact propagator D satisfies the variational equation

$$\frac{\delta\Gamma[\phi_0, \rho_0, D]}{\delta D} = 0. \quad (17)$$

Using Eq. (17) and $\Pi = D^{-1} - D_0^{-1}$, where Π is the exact self-energy, the variational equation can be written as

$$\Pi(D) = 2\frac{\delta\Phi[D]}{\delta D}. \quad (18)$$

The diagrams that contribute to the self-energy matrix $\Pi(\omega, p)$ are obtained by cutting a line in the vacuum diagrams.

In Ref. [21], the diagrams contributing to $\Phi[D]$ were classified according to which order in the $1/N$ expansion they contribute. The various terms can be written in terms of $O(N)$ invariants such as $\text{Tr}(D^n)$ and $\text{Tr}(\phi_0^2 D^n)$, where n is an integer. At leading order in $1/N$ the diagrams that contribute are shown in Fig. 1. This diagram can be written as an $O(N)$ -invariant term, $\Phi_{\text{LO}} \sim (\text{Tr} D)^2$. The vertex gives a factor of $1/N$, while each trace yields a factor of N . It is therefore a leading-order contribution.

The diagram that contributes to the self-energy Π to leading in $1/N$ is shown in Fig. 2. The particle propagating in the loop is one of the $2N - 3$ particles with tree-level masses given by Eq. (13).




FIG. 1: Feynman diagram contributing at leading order in $1/N$ to $\Phi[D]$.




FIG. 2: Feynman diagram contributing at leading order in $1/N$ to the self-energy $\Pi(\omega, p)$.

Their inverse propagator to leading order in $1/N$ can be written as

$$D^{-1}(\omega_n, p) = P^2 + m_4^2 + \Pi_{\text{LO}}(D), \quad (19)$$

where $P^2 = \omega_n^2 + p^2$ and $\Pi_{\text{LO}}(D)$ is the tadpole in Fig. (2) calculated with the full propagator D . Since $\Pi_{\text{LO}}(D)$ is independent of the external momentum, the propagator can be parametrized as

$$D^{-1}(\omega_n, p) = P^2 + M^2, \quad (20)$$

where M is a constant. The leading order self energy can now be written as

$$\Pi_{\text{LO}} = \lambda \oint_Q \frac{1}{Q^2 + M^2}. \quad (21)$$

Using the expression for Π_{LO} and the two expressions (19) and (20) for the inverse propagator, we obtain a gap equation for M :

$$M^2 = m_4^2 + \lambda \oint_Q \frac{1}{Q^2 + M^2}. \quad (22)$$

The sum-integral is defined by

$$\oint_Q \equiv \left(\frac{e^{\gamma} \Lambda^2}{4\pi} \right)^\epsilon T \sum_{q_0=2\pi n T} \int_p \frac{d^d q}{(2\pi)^d}, \quad (23)$$

where $d = 3 - 2\epsilon$ and Λ is the renormalization scale associated with dimensional regularization. The integral over 3-momentum will be calculated using dimensional regularization. Inserting the parametrization (20) of D

into the 2PI effective potential (14) and differentiating with respect to ϕ_0 and ρ_0 , the equations (15) and (16) can be written as

$$0 = m^2 \phi_0 - H + \frac{\lambda}{2N} \phi_0 (\phi_0^2 + \rho_0^2) + \lambda \phi_0 \oint_Q \frac{1}{Q^2 + M^2}, \quad (24)$$

$$0 = (m^2 - \mu_I^2) \rho_0 + \frac{\lambda}{2N} \rho_0 (\phi_0^2 + \rho_0^2) + \lambda \rho_0 \oint_Q \frac{1}{Q^2 + M^2}. \quad (25)$$

Combining the gap equation (22) and Eq. (24) gives

$$H = M^2 \phi_0. \quad (26)$$

Similarly, combining the gap equation (22) and Eq. (25) gives

$$(M^2 - \mu^2) \rho_0 = 0. \quad (27)$$

We will study the various phases in the following subsections.

We next discuss the renormalization of the gap equation (22), which can be done by an iterative procedure given in Ref. [25] (see also Ref. [41]). By writing $M^2 = m_4^2 + \Pi_{\text{LO}}$ and expanding the right-hand side of Eq. (22) in powers of Π_{LO} , it is clear that it resums the perturbative daisy and superdaisy diagrams:

$$\begin{aligned} M^2 - m_4^2 &= \lambda \oint_Q \frac{1}{Q^2 + m_4^2 + \Pi_{\text{LO}}} \\ &= \lambda \oint_Q \frac{1}{Q^2 + m_4^2} - \Pi_{\text{LO}} \lambda \oint_Q \frac{1}{(Q^2 + m_4^2)^2} \\ &\quad + \Pi_{\text{LO}}^2 \lambda \oint_Q \frac{1}{(Q^2 + m_4^2)^3} + \dots \end{aligned} \quad (28)$$




FIG. 3: Diagrammatic interpretation of the gap equation (22) for Π_{LO} .

This is illustrated diagrammatically in Fig 3. Renormalization is carried by making the substitutions

$$m^2 \rightarrow m^2 + \sum_{n=1}^{\infty} \delta m_n^2, \quad (29)$$

$$\lambda \rightarrow \lambda + \sum_{n=1}^{\infty} \delta \lambda_n, \quad (30)$$

where δm_n^2 and $\delta \lambda_n$ are counterterms of order λ^n and λ^{n+1} , respectively. Similarly, we write the leading order self-energy in a power series in λ :

$$\Pi_{\text{LO}} = \Pi_{\text{LO},1} + \Pi_{\text{LO},2} + \dots, \quad (31)$$

where the subscript indicates the power of λ . Inserting Eqs. (29)–(31) into equation (28), we can determine $\Pi_{\text{LO},n}$, δm_n^2 and $\delta \lambda_n$ by iteration. The first iteration of the gap equation yields

$$\Pi_{\text{LO},1} = \lambda \oint_Q \frac{1}{Q^2 + m_4^2} + \delta m_1^2 + \frac{\delta \lambda_1}{2N} (\phi_0^2 + \rho_0^2), \quad (32)$$

where we have used that $\Pi_{\text{LO}} = M^2 - m_4^2$. The next step is to calculate the sum-integral in Eq. (32). The sum over Matsubara frequencies can be expressed as a contour integral in the complex energy plane. By performing this integral, we obtain

$$\oint_Q \frac{1}{Q^2 + M^2} = \int_q \frac{d^d q}{(2\pi)^d} \frac{1}{\sqrt{Q^2 + M^2}} \times \left[\frac{1}{2} + \frac{1}{e^{\sqrt{Q^2 + M^2}/T} - 1} \right]. \quad (33)$$

The first term is divergent in the ultraviolet. Calculating it with dimensional regularization and expanding in ϵ through order ϵ^0 , we obtain

$$\oint_Q \frac{1}{Q^2 + M^2} = -\frac{M^2}{16\pi^2} \left(\frac{\Lambda^2}{M^2} \right)^\epsilon \left[\frac{1}{\epsilon} + 1 \right] + \frac{1}{2\pi^2} \int_0^\infty \frac{dq q^2}{\sqrt{q^2 + M^2}} \frac{1}{e^{\sqrt{q^2 + M^2}/T} - 1}. \quad (34)$$

The logarithmic divergence in the sum-integral in Eq. (22) shows up as a pole in ϵ , while the quadratic divergence is set to zero in dimensional regularization. Inserting the sum-integral (34) into Eq. (32), we obtain

$$\Pi_{\text{LO},1} = -\frac{\lambda m_4^2}{16\pi^2 \epsilon} + \delta m_1^2 + \frac{\delta \lambda_1}{2N} (\phi_0^2 + \rho_0^2)$$

$$+ \text{finite terms}. \quad (35)$$

The divergences are removed by the counterterms

$$\delta m_1^2 = \frac{m^2 \lambda}{16\pi^2 \epsilon}, \quad \delta \lambda_1 = \frac{\lambda^2}{16\pi^2 \epsilon}. \quad (36)$$

This renormalization procedure can be carried out iteratively to all orders in λ and this determines the counterterms δm_n^2 and $\delta \lambda_n$ for $n > 1$. The counterterms δm_n^2 and $\delta \lambda_n$ are [25]

$$\delta m_n^2 = \frac{m^2 \lambda^n}{(16\pi^2 \epsilon)^n}, \quad \delta \lambda_n = \frac{\lambda^{n+1}}{(16\pi^2 \epsilon)^n}. \quad (37)$$

Summing the counterterms to all orders, Eqs. (29) and (30) can be written as

$$m^2 \rightarrow \frac{m^2}{1 - \frac{\lambda}{16\pi^3 \epsilon}}, \quad \lambda \rightarrow \frac{\lambda}{1 - \frac{\lambda}{16\pi^3 \epsilon}}. \quad (38)$$

Returning to the nonperturbative gap equation (22), it can be rewritten as

$$\begin{aligned} \frac{M^2}{\lambda} &= \frac{m^2}{\lambda} + \frac{1}{2N} (\phi_0^2 + \rho_0^2) + \oint_Q \frac{1}{Q^2 + M^2} \\ &= \frac{m^2}{\lambda} + \frac{1}{2N} (\phi_0^2 + \rho_0^2) - \frac{M^2}{16\pi^2} \left(\frac{\Lambda^2}{M^2} \right)^\epsilon \left[\frac{1}{\epsilon} + 1 \right] + \frac{1}{2\pi^2} \int_0^\infty \frac{dq q^2}{\sqrt{q^2 + M^2}} \frac{1}{e^{\sqrt{q^2 + M^2}/T} - 1}. \end{aligned} \quad (39)$$

Making the substitutions (38), the renormalized gap

equation becomes

$$M^2 = m^2 + \frac{\lambda}{2N} (\phi_0^2 + \rho_0^2) - \frac{\lambda M^2}{16\pi^2} \left(\ln \frac{\Lambda^2}{M^2} + 1 \right)$$

$$+ \frac{\lambda}{2\pi^2} \int_0^\infty \frac{dq q^2}{\sqrt{q^2 + M^2}} \frac{1}{e^{\sqrt{q^2 + M^2}/T} - 1} . \quad (40)$$

We finally discuss the renormalization of the effective potential in the broken phase. In the large- N limit, $\Phi[D]$ is given by

$$\Phi[D] = \frac{1}{2} \lambda N \left(\oint_Q \frac{1}{P^2 + M^2} \right)^2 . \quad (41)$$

Using the expression (41) and inserting the gap equation (22) into the expression for the effective potential (14), we obtain

$$\Gamma[\phi_0, \rho_0, D] = \frac{N}{2\lambda} (M^4 - m^4) - \frac{1}{2} \mu_I^2 \rho_0^2 - H \phi_0 - N M^2 \oint_Q \frac{1}{P^2 + M^2} + N \oint_Q \ln(P^2 + M^2) . \quad (42)$$

The first sum-integral is given by Eq. (34), while the second can be found by integrating the first with respect to M^2 and reads

$$\oint_Q \ln(P^2 + M^2) = -\frac{M^4}{32\pi^2} \left(\frac{\Lambda^2}{M^2} \right)^\epsilon \left[\frac{1}{\epsilon} + \frac{3}{2} \right] + \frac{T}{\pi^2} \int_0^\infty dq q^2 \ln \left(1 - e^{-\sqrt{q^2 + M^2}/T} \right) . \quad (43)$$

Inserting the expressions for the sum-integrals into (42) and making the substitutions (38), we obtain the renormalized effective potential

$$\Gamma[\phi_0, \rho_0, D] = \frac{N}{2\lambda} (M^4 - m^4) - \frac{1}{2} \mu_I^2 \rho_0^2 - H \phi_0 + \frac{N M^4}{32\pi^2} \left[\ln \frac{\Lambda^2}{M^2} + \frac{1}{2} \right] - \frac{N M^2}{2\pi^2} \int_0^\infty \frac{dq q^2}{\sqrt{q^2 + M^2}} \frac{1}{e^{\sqrt{q^2 + M^2}/T} - 1} + \frac{N T}{\pi^2} \int_0^\infty dq q^2 \ln \left(1 - e^{-\sqrt{q^2 + M^2}/T} \right) , \quad (44)$$

where we have omitted a temperature-independent divergence proportional to m^4 . This corresponds to adding a vacuum counterterm $\Delta\mathcal{E}$ to the effective potential. We also note that the effective potential can be renormalized in a similar manner in the symmetric phase.

A. Chiral limit

In this subsection, we discuss the solutions to the gap equations and the corresponding phases in the chiral limit, i.e. when $H = 0$.

First assume that the charged pion condensate ρ_0 is nonzero. Equation (27) then immediately gives

$$M^2 = \mu_I^2 . \quad (45)$$

Equation (26) then implies that the chiral condensate vanishes:

$$\phi_0 = 0 . \quad (46)$$

Using Eqs. (25), (45) and (46), the pion condensate reads

$$\rho_0^2 = \frac{2N}{\lambda} (\mu_I^2 - m^2) - 2N \oint_Q \frac{1}{Q^2 + \mu_I^2} . \quad (47)$$

The pole in ϵ on the right-hand side of this equation is removed by making the substitutions (38) in analogy with the renormalization of the gap equation (22). After renormalization, the gap equation (47) can be written as

$$\rho_0^2 = \frac{2N}{\lambda} (\mu_I^2 - m^2) - \frac{N \mu_I^2}{8\pi^2} \left(\ln \frac{\Lambda^2}{\mu_I^2} + 1 \right) - \frac{N}{\pi^2} \int_0^\infty \frac{dq q^2}{\sqrt{q^2 + \mu_I^2}} \frac{1}{e^{\sqrt{q^2 + \mu_I^2}/T} - 1} . \quad (48)$$

As the temperature T increases, Eq. (48) shows that ρ_0 goes to zero continuously and so the phase transition is second order. The critical temperature T_c is given by the temperature for which the right-hand side of Eq. (48) vanishes. In the weak-coupling limit, where $\lambda \ll 1$, we can obtain an analytic expression for T_c . In that case, μ_I can be neglected in the sum-integral in Eq. (47) and its value is $NT^2/6$. This yields the critical temperature

$$T_c^2 = \frac{12}{\lambda} (\mu_I^2 - m^2) , \quad (49)$$

which is in agreement with the calculations of Kapusta [42].

The dispersion relations are obtained by analytic continuation to Minkowski space, $\omega_n \rightarrow i\omega$, and then solving the equation $\det D(\omega, p) = 0$. For $T < T_c$, the dispersion relations are

$$\omega_1(p) = \sqrt{p^2 + \mu_I^2} , \quad (50)$$

$$\omega_{2,3}(p) = \sqrt{p^2 + 2\mu_I^2 + \frac{1}{2}(M_2^2 + M_3^2) \pm \sqrt{(M_2^2 - M_3^2)^2 + 8\mu_I^2(2\mu_I^2 + M_2^2 + M_3^2) + 4\mu_I^2 p^2}}, \quad (51)$$

$$\omega_4(p) = \sqrt{p^2 + \mu_I^2}, \quad (52)$$

where $M_i^2 = m_i^2 + \Pi_{\text{LO}}$. Note that the σ and π^0 are identified with the original fields ϕ_1 and ϕ_4 , and therefore with $\omega_1(p)$ and $\omega_4(p)$. On the other hand, the charged pions π^- and π^+ are given by linear combinations of ϕ_2 and ϕ_3 and are identified by $\omega_2(p)$ and $\omega_3(p)$. In the minimum of the effective potential (14), we have $M_2^2 = 2(\mu_I^2 - \tilde{m}^2)$ and $M_3^2 = 0$, where $\tilde{m}^2 = m^2 + \Pi_{\text{LO}}$. The dispersion relations in Eq. (51) then reduce to

$$\omega_{2,3}(p) = \sqrt{p^2 + 3\mu_I^2 - \tilde{m}^2 \pm \sqrt{(3\mu_I^2 - \tilde{m}^2)^2 + 4\mu_I^2 p^2}}. \quad (53)$$

From this equation, we see that the π^+ is a massless mode, which in the long-wavelength limit behaves as

$$\omega_3(p) = \sqrt{\frac{\mu_I^2 - \tilde{m}^2}{3\mu_I^2 - \tilde{m}^2}} p. \quad (54)$$

This is consistent with the fact that the original $O(2) \times O(2)$ symmetry is broken down to $O(2)$ by the pion condensate. The fact that the Goldstone mode is behaving linearly with p makes the system superfluid as the Landau criterion is satisfied [43].

In the normal phase, where $\phi_0 = \rho_0 = 0$, the dispersion relations are

$$\omega_1(p) = \sqrt{p^2 + M^2}, \quad (55)$$

$$\omega_{2,3}(p) = \sqrt{p^2 + M^2} \pm \mu_I, \quad (56)$$

$$\omega_4(p) = \sqrt{p^2 + M^2}, \quad (57)$$

where M is determined selfconsistently by solving the

gap equation (40) evaluated at $\phi_0 = \rho_0 = 0$:

$$M^2 = m^2 - \frac{\lambda M^2}{16\pi^2} \left(\ln \frac{\Lambda^2}{M^2} + 1 \right) + \frac{\lambda}{2\pi^2} \int_0^\infty \frac{dq q^2}{\sqrt{q^2 + M^2}} \frac{1}{e^{\sqrt{q^2 + M^2}/T} - 1}. \quad (58)$$

B. Physical point

We next discuss the case where $H \neq 0$. The solution to Eq. (26) yields

$$\phi_0 = \frac{H}{\mu_I^2}. \quad (59)$$

Note that ϕ_0 is nonzero and independent of the temperature. Assuming a nonzero pion condensate, the mass M again is given by Eq. (45). Using Eqs. (45) and (59), the gap equation (25) can be written as

$$\rho_0^2 = \frac{2N}{\lambda} (\mu_I^2 - m^2) - \frac{H^2}{\mu_I^4} - 2N \oint_Q \frac{1}{Q^2 + \mu_I^2}. \quad (60)$$

The gap equation is renormalized in the same way as the gap equation in the chiral limit.

The expressions for the dispersion relations are very involved and they are not particularly illuminating. We therefore restrict ourselves to list the quasiparticle masses. These are found by solving the equation $\det D(\omega, p = 0) = 0$ and read

$$\omega_1(p = 0) = M_1 = 0, \quad (61)$$

$$\omega_{2,3}(p = 0) = M_{2,3} = \sqrt{\frac{7}{2}\mu_I^2 - \tilde{m}^2 \pm \frac{1}{2}\sqrt{(5\mu_I^2 - 2\tilde{m}^2)^2 - 12\frac{\lambda}{N}\frac{H^2}{\mu_I^2}}}, \quad (62)$$

$$\omega_4(p = 0) = M_4 = \mu_I, \quad (63)$$

where $\tilde{m}^2 = m^2 + \Pi_{\text{LO}}$. The π^+ is again a massless Goldstone mode. By expanding the equation for the

dispersion relations around $p = 0$, it can be shown that this mode is linear in p and hence the system is super-

fluid. This mode reflects the breaking of the residual $O(2)$ symmetry and is a conventional Goldstone mode.

We next discuss the system in the normal phase, where $\rho_0 = 0$. Eliminating ϕ_0 from the gap equation (24), one finds

$$\frac{2N}{\lambda} (M^2 - m^2) = \frac{H^2}{M^4} + 2N \oint_Q \frac{1}{Q^2 + M^2}. \quad (64)$$

This equation is renormalized in the usual manner. The renormalized gap equation determines the mass M as a function of temperature. We see that the equations (60) and (64) match onto each other continuously at the phase boundary, where $M = \mu_I$ as they should. The dispersion relations in the symmetric phase

$$\omega_1(p) = \sqrt{p^2 + M^2 + \frac{\lambda}{N} \frac{H^2}{M^4}}, \quad (65)$$

$$\omega_{2,3}(p) = \sqrt{p^2 + M^2} \pm \mu_I, \quad (66)$$

$$\omega_4(p) = \sqrt{p^2 + M^2}, \quad (67)$$

Note that all four modes are massive as the $O(2)$ symmetry has been restored.

C. Numerical results

We next discuss the determination of the parameters in the Lagrangian (1). In the vacuum, we have $\rho_0 = \mu_I = 0$ and $f_\pi = \phi_0$. At the tree-level, we have

$$m_\sigma^2 = m_1^2, \quad (68)$$

$$m_\pi^2 = m_2^2, \quad (69)$$

$$H = f_\pi m_\pi^2. \quad (70)$$

The mass parameter m^2 and the coupling constant λ can then be expressed in terms of the observable masses of the sigma and the pion:

$$m^2 = -\frac{1}{2}(m_\sigma^2 - 3m_\pi^2), \quad (71)$$

$$\lambda = N \frac{(m_\sigma^2 - m_\pi^2)}{f_\pi^2}. \quad (72)$$

Eqs. (70)–(72) can now be used to determine the constants H , m^2 , and λ , given the values $m_\sigma = 600\text{MeV}$, $m_\pi = 139\text{MeV}$, and $f_\pi = 93\text{MeV}$. In the chiral limit, we find $m^2 = -(300\text{MeV})^2$ and $\lambda = 83.2466$. At the physical point, we obtain $m^2 = -(230.5\text{MeV})^2$ and $\lambda = 78.7788$. Once we take into account quantum fluctuations, Eqs. (68) and (69) are modified. After renormalization, we find

$$m_\sigma^2 = m_2^2 - \frac{\lambda m_\pi^2}{16\pi^2} \ln \frac{\Lambda^2 e}{m_\pi^2}. \quad (73)$$

$$m_\pi^2 = m_1^2 - \frac{\lambda m_\pi^2}{16\pi^2} \ln \frac{\Lambda^2 e}{m_\pi^2}. \quad (74)$$

Note that the renormalization scale Λ is a free parameter and in the chiral limit, it does not enter our expressions. At the physical point we chose $\Lambda^2 = m_\pi^2/e$, so that the tree-level relations still hold without having to change the parameters m^2 and λ .

In Fig. 4, we show the medium dependent meson masses at the physical point as functions of the isospin chemical potential at zero temperature. For $\mu_I > \mu_c = 139\text{ MeV}$, they are given by Eqs. (61)–(63), while for $\mu_I < \mu_c$, they are given by Eqs. (65)–(67) with $p = 0$. For $\mu_I > \mu_c$, the mass of π^+ vanishes.




FIG. 4: Pion and sigma masses at the physical point as functions of the isospin chemical potential μ_I and $T = 0$.

In Fig. 5, we show the pion condensate as a function of temperature normalized to the condensate at $T = 0$. In both the chiral limit and at the physical point, the pion condensate breaks an $O(2)$ symmetry. From universality arguments, we know this transition is of second order, which can be seen from Fig. (5). The critical exponent $\nu = 1/2$, which is expected from a mean-field calculation. See Ref. [44] for a NLO calculation of the critical exponents using the 2PI formalism.

In Fig. 6, we show the phase diagram for pion condensation. The solid line corresponds to the chiral limit, while the dashed line corresponds to the physical point. The two curves meet at large μ_I since the temperature-independent value of H becomes irrelevant.

In Fig. 7, we show the meson masses as functions of temperature for fixed isospin chemical potential $\mu_I = 200\text{ MeV}$. For $T < T_c$, they are given by Eqs. (61)–(63), while for $T > T_c$, they are given by Eqs. (65)–(67) with $p = 0$. The critical temperature is $T_c = 223\text{ MeV}$. Note that one of the pions is massless below the critical temperature




FIG. 5: Charged pion condensate as a function of temperature normalized to the condensate at $T = 0$. The upper curve is the chiral limit and the lower curve corresponds to the physical point.




FIG. 6: Phase diagram for pion condensation. The upper curve is the chiral limit and the lower curve corresponds to the physical point.

IV. KAONS

In the color-flavor locked phase of high baryon-density QCD, the lightest excitations are the four charged and

neutral kaons, $K^\pm$, K^0 and $\bar{K}^0$. The low-energy effective theory is described by the Lagrangian (1) with $H = 0$ and $N = 2$ [5]. In this section, we discuss kaon condensation in the presence of chemical potentials μ_Y for hypercharge Y and μ_I for isospin. The kaons can be written as a complex doublet, $(\Phi_1, \Phi_2) = (K^0, K^+)$. We then have the following relation between the various




FIG. 7: Meson masses as functions of temperature at the physical point. The isospin chemical potential is $\mu_I = 200$ MeV. $T_c = 223$ MeV.

chemical potentials:

$$\mu_I = \frac{1}{2}(\mu_2 - \mu_1), \quad (75)$$

$$\mu_Y = \mu_1 + \mu_2. \quad (76)$$

We first write the complex fields Φ_1 and Φ_2 in terms of the charged and neutral kaon condensates ϕ_0 and ρ_0 and quantum fluctuating fields $\phi_1 - \phi_4$:

$$\Phi_1 = \frac{1}{\sqrt{2}}(\phi_0 + \phi_1 + i\phi_2), \quad (77)$$

$$\Phi_2 = \frac{1}{\sqrt{2}}(\rho_0 + \phi_3 + i\phi_4). \quad (78)$$

The tree-level propagator is

$$D_0^{-1}(\omega_n, p) = \begin{pmatrix} \omega_n^2 + p^2 + m_1^2 & -2\mu_1\omega_n & \frac{\lambda}{N}\phi_0\rho_0 & 0 & 0 & \dots \\ 2\mu_1\omega_n & \omega_n^2 + p^2 + m_2^2 & 0 & 0 & 0 & \dots \\ \frac{\lambda}{N}\phi_0\rho_0 & 0 & \omega_n^2 + p^2 + m_3^2 & -2\mu_2\omega_n & 0 & \dots \\ 0 & 0 & 2\mu_2\omega_n & \omega_n^2 + p^2 + m_4^2 & 0 & \dots \\ \dots & \dots & \dots & \dots & \omega_n^2 + p^2 + m_5^2 & \dots \end{pmatrix}, \quad (79)$$

where the tree-level masses are

$$m_1^2 = -\mu_1^2 + m^2 + \frac{3\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2, \quad (80)$$

$$m_2^2 = -\mu_1^2 + m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2, \quad (81)$$

$$m_3^2 = -\mu_2^2 + m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{3\lambda}{2N}\rho_0^2, \quad (82)$$

$$m_4^2 = -\mu_2^2 + m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2, \quad (83)$$

$$m_5^2 = m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2. \quad (84)$$

Note that in the remainder of this section, the mass parameter m^2 is positive. The 2PI effective potential is given by

$$\begin{aligned} \Gamma[\phi_0, \rho, D] &= \frac{1}{2}(m^2 - \mu_1^2)\phi_0^2 + \frac{1}{2}(m^2 - \mu_2^2)\rho_0^2 \\ &+ \frac{\lambda}{8N}(\phi_0^2 + \rho_0^2)^2 + \frac{1}{2}\text{Tr} \ln D^{-1} \\ &- \frac{1}{2}\text{Tr} D_0^{-1}D + \Phi[D]. \end{aligned} \quad (85)$$

To leading order in N , the contribution to $\Phi[D]$ is again given by the diagrams in Fig. 1. The stationarity conditions (15)–(16) for the effective potential are

$$\begin{aligned} 0 &= (m^2 - \mu_1^2)\phi_0 + \frac{\lambda}{2N}\phi_0(\phi_0^2 + \rho_0^2) \\ &+ \lambda\phi_0 \oint_Q \frac{1}{Q^2 + M^2}, \\ 0 &= (m^2 - \mu_2^2)\rho_0 + \frac{\lambda}{2N}\rho_0(\phi_0^2 + \rho_0^2) \end{aligned} \quad (86)$$

$$+ \lambda\rho_0 \oint_Q \frac{1}{Q^2 + M^2}. \quad (87)$$

The self-energy to leading order in $1/N$ is again diagonal and contributing diagram is shown in Fig. 2. The expression is given by Eq. (22). Using Eq. (86) for the minimum of the effective potential, we obtain

$$M^2 = \mu_1^2. \quad (88)$$

We first assume that $\mu_1 = \mu_2$. In this case, the Lagrangian (1) has an extended $SU(2) \times U(1)$ symmetry instead of $O(2) \times O(2)$. If $\mu_1^2 > m^2$ this symmetry is broken down to $U(1)$. We have complete symmetry between ϕ_0 and ρ_0 and it is then possible to rotate e.g. the ρ_0 condensate away. If $\mu_1 > \mu_2$ (to be specific), $\rho_0 = 0$ and $\phi_0 \neq 0$ only if $\mu_1^2 > m^2$. In this case the $O(2) \times O(2)$ symmetry is spontaneously broken down to $O(2)$ by the condensate.

In analogy with Eq. (24), the kaon condensate ϕ_0 satisfies the gap equation

$$\phi_0^2 = \frac{2N}{\lambda}(\mu_1^2 - m^2) - 2 \oint_Q \frac{1}{Q^2 + M^2}. \quad (89)$$

The phase transition is again of second order. In the weak-coupling limit, an analytic expression for T_c analogous to Eq. (49) is obtained by the substitution $\mu_I \rightarrow \mu_1$.

In the broken phase, the dispersion relations can be written as

$$\omega_{1,2}(p) = \sqrt{p^2 + 2\mu_1^2 + \frac{1}{2}(M_1^2 + M_2^2) \pm \sqrt{(M_1^2 - M_2^2)^2 + 8\mu_1^2(2\mu_1^2 + M_1^2 + M_2^2) + 4\mu_1^2 p^2}}, \quad (90)$$

$$\omega_{3,4}(p) = \sqrt{p^2 + \mu_1^2 \pm \mu_2}, \quad (91)$$

$$\omega_5(p) = \sqrt{p^2 + \mu_1^2}, \quad (92)$$

where $M_i^2 = m_i^2 + \Pi_{\text{LO}}$. The neutral kaons $\bar{K}^0$ and K^0 are identified with the linear combinations of the fields ϕ_1 and ϕ_2 , and therefore with $\omega_1(p)$ and $\omega_2(p)$. Similarly, the charged kaons K^- and K^+ are given by linear combinations of ϕ_3 and ϕ_4 and are identified with $\omega_3(p)$ and $\omega_4(p)$.

At the minimum of the effective potential, we have $M_1^2 = 2(\mu_1^2 - \tilde{m}^2)$ and $M_2^2 = 0$, where $\tilde{m}^2 = m^2 + \Pi_{\text{LO}}$. Eq. (90) then reduce to

$$\omega_{1,2}(p) = \sqrt{p^2 + 3\mu_1^2 - \tilde{m}^2 \pm \sqrt{(3\mu_1^2 - \tilde{m}^2)^2 + 4p^2\mu_1^2}}. \quad (93)$$

This shows that $\omega_2(p)$ and thus K^0 is a massless excitation which is linear for small p (cf. (54)) and has the quantum numbers of K^0 . In the case where $\mu_1 \neq \mu_2$, this is the only Goldstone boson which corresponds to the broken $O(2)$ symmetry. In the case where $\mu_1 = \mu_2$, $\omega_4(p)$ is another massless modes, which is quadratic for small p , which has the quantum numbers of K^+ . Thus there are two massless modes despite the fact that there are 3 broken generators [45, 46, 47]. The splitting between K^+ and K^- is due to the fact that the chemical potentials break the discrete symmetries C , CP , and CPT [47]. This result is in agreement with the counting rules derived by Nielsen and Chadha [48], which state that the massless modes with a quadratic dispersion relation must be counted twice. Finally, we notice that only in the case where $\mu_1 \neq \mu_2$, the condensate implies that the system is superfluid. Whenever the Goldstone mode with a quadratic dispersion relation is present, the Landau criterion for superfluidity can never be satisfied [46, 47].

We next discuss the system in the normal phase, where $\phi_0 = 0$. The dispersion relations reduce to

$$\omega_{1,2}(p) = \sqrt{p^2 + M^2} \pm \mu_1, \quad (94)$$

$$\omega_{3,4}(p) = \sqrt{p^2 + M^2} \pm \mu_2, \quad (95)$$

$$\omega_5(p) = \sqrt{p^2 + M^2}, \quad (96)$$

where M^2 again satisfies the gap equation (58). Since the symmetry is restored, all excitations are gapped.

In Fig. 8, we show the quasiparticle masses as a function of μ_1 at zero temperature. These are given by Eqs. (90)-(92) with $p = 0$ in the broken phase and by Eqs. (94)-(96) with $p = 0$ in the symmetric phase. In the broken phase, the neutral kaon K^0 is a massless Goldstone mode. We have set $\mu_2 = \mu_1/2$ and chosen the renormalization scale $\Lambda = M$ in the symmetric phase. The phase diagram is the same as in the pion case with $H \neq 0$.




FIG. 8: Kaon masses at zero temperature as functions of the chemical potential μ_1 , where $\mu_1 > \mu_2$.

V. SUMMARY

In the present paper, we have calculated to leading order in the 2PI-1/ N expansion finite density and finite-temperature effects in interacting Bose systems with pion and kaon condensation. In particular, we have obtained the phase diagrams and quasiparticle masses. Depending on the magnitude of the chemical potentials, the number and nature of the Goldstone modes vary. In all cases, our results are in agreement with general theorems on the occurrence of massless excitations [45, 48, 49]. This is in contrast with e.g. the Hartree approximation, where the Goldstone theorem is not satisfied. Furthermore, we have shown that it is possible to renormalize the theory in a medium-independent way by eliminating the divergences through a redefinition of the mass and coupling constant.

The present work can be extended by calculating the 1/ N corrections either in the context of the standard effective action or by using the 2PI effective-action formalism. Using the conventional 1PI effective action, the calculations would be similar to those recently carried out in Refs. [21, 22]. Calculations within the 2PI effective action will be more difficult as the resulting equations form a set of coupled integro-differential equations. These equations must be solved numerically, which is nontrivial. However, progress has been made recently [14] in massive scalar field theory with vanishing chemical potential.

Acknowledgment

The author would like to thank D. Boer and H. J. Warringa for useful discussion and suggestions.

-
- [1] L. P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation*, Oxford University Press, Oxford, UK (2003).
 - [2] J. O. Andersen, Rev. Mod. Phys. **76**, 599 (2004).
 - [3] P. F. Bedaque and T. Schäfer, Nucl. Phys. A **697**, 802 (2002).
 - [4] D. B. Kaplan and S. Reddy, Phys. Rev. D **65**, 054042 (2002).
 - [5] M. G. Alford, K. Rajagopal, and F. Wilczek, Nucl. Phys. **B537**, 443 (1999).
 - [6] J.-P. Blaizot, E. Iancu, and A. K. Rebhan, hep-ph/0303185.
 - [7] U. Kraemmer and A. K. Rebhan, Rept. Prog. Phys. **67** (2004).
 - [8] J. O. Andersen and M. Strickland, Ann. Phys. **317**, 281 (2005).
 - [9] F. Karsch, A. Patkós, and P. Petreczky, Phys. Lett. B **401**, 69 (1997).
 - [10] J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. Lett. **83** 2139 (1999), Phys. Rev. D **66**, 085016 (2002).
 - [11] J.-P. Blaizot, E. Iancu, and A. Rebhan, Phys. Lett. B **470**, 181, (1999); Phys. Rev. Lett. **83**, 2906 (1999); Phys. Rev. D **63**, 065003 (2001).
 - [12] E. Braaten and E. Petigirard, Phys. Rev. D **65**, 041701, (2002); ibid 085039 (2002).
 - [13] J. O. Andersen and M. Strickland, Phys. Rev. D **71**, 025011 (2005).
 - [14] J. Berges, Sz. Borsanyi, U. Reinosa, and J. Serreau, Phys. Rev. D **71**, 105004 (2005).
 - [15] B. A. Freedman and L. D. McLerran, Phys. Rev. D **16**, 1169 (1977).
 - [16] V. Baluni, Phys. Rev. D **17**, 2092 (1978).
 - [17] E. S. Fraga, R. D. Pisarski, and J. Schaffner-Bielich, Phys. Rev. D **63**, 121702 (2001).
 - [18] A. Ipp, A. Gerhold, and A. Rebhan, Phys. Rev. D **69**, 011901 (2004).
 - [19] A. Ipp, K. Kajantie, A. Rebhan, and Vuorinen, Phys. Rev. D **74**, 045016 (2006).
 - [20] H. Meyer-Ortmanns, Rev. Mod. Phys. **68**, 473 (1996).
 - [21] G. Aarts, D. Ahrensmeier, R. Baier, J. Berges, and J. Serreau, Phys. Rev. D **66**, 045008, (2002).
 - [22] J. O. Andersen, D. Boer, and H. J. Warringa, Phys. Rev. D **69**, 076006 (2004); ibid **70**, 116007 (2004).
 - [23] J. T. Lenaghan and D. H. Rischke, J. Phys. **G26**, 431 (2000).
 - [24] H. van Hees, J. Knoll, Phys. Rev. D **65**, 025010 (2002); ibid 65, 105005 (2002); 65, 025028 (2002).
 - [25] J.-P. Blaizot, E. Iancu and U. Reinosa, Phys. Lett. B **568**, 160 (2003), Nucl. Phys. A **736**, 149 (2004).
 - [26] J. Berges, Sz. Borzányi, U. Reinosa, and J. Serreau, Ann. Phys. **320**, 344 (2005).
 - [27] D. Page and S. Reddy, astro-ph/0608360.
 - [28] D. T. Son and M. Stephanov, Phys. Rev. Lett. **86**, 592 (2001).
 - [29] K. Splittorff, D. T. Son and M. Stephanov, Phys. Rev. D **64** 016003 (2001).
 - [30] J. B. Kogut and D. Toublan, Phys. Rev. D **64** 034007 (2001).
 - [31] M. Loewe and C. Villavicencio, Phys. Rev. D **67**, 074034 (2003); ibid D **70**, 074005 (2004).
 - [32] J. B. Kogut and D. K. Sinclair, Phys. Rev. D **64** 014508 (2002); ibid D **66** 34505 (2002).
 - [33] S. Gupta, hep-lat/0202005.
 - [34] A. Barducci, R. Casalbuoni, G. Pettini, and L. Ravagli, Phys. Lett. **B564**, 217 (2003). D **71**, 094001 (2005).
 - [35] A. Jakovac, A. Patkos, Zs. Szep, and P. Szepfalusy, Phys. Lett. **B582**, 179 (2004).
 - [36] A. Barducci, R. Casalbuoni, G. Pettini, and L. Ravagli, Phys. Rev. D **69**, 096004 (2004).
 - [37] L. He, M. Jin, and P. Zhuang, Phys. Rev. D **71**, 116001 (2005).
 - [38] D. Ebert and K. G. Klimenko, J. Phys. **G32**, 599 (2006); Eur. Phys. J. C **46**, 771 (2006).
 - [39] H. Mao, N. Petropoulos, and W.-Q. Zhao, J. Phys. **G32**, 2187 (2006).
 - [40] H. E. Haber and H. A. Weldon, Phys. Rev. D **25**, 502, (1982).
 - [41] A. Patkos and Zs. Szep, Phys. Lett. **B642**, 384 (2006).
 - [42] J. I. Kapusta, Phys. Rev. D **24**, 426, (1981).
 - [43] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics, 2nd edition*, Pergamon New York (1987).
 - [44] M. Alford, J. Berges, and J. M. Cheyne, Phys. Rev. D **70**, 125002 (2004).
 - [45] T. Schäfer, D. T. Son, M. A. Stephanov, D. Toublan, and J. J. M. Verbaarschot, Phys. Lett. **B522**, 67 (2001).
 - [46] V. A. Miransky and I. A. Shovkovy, Phys. Rev. Lett. **88**, 111601 (2002).
 - [47] V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Mod. Phys. Lett. **A19**, 1341 (2004).
 - [48] H. B. Nielsen and S. Chadha, Nucl. Phys. **B105**, 445 (1976).
 - [49] T. Brauner, Phys. Rev. D **72**, 076002 (2005); PhD Thesis, Nuclear Physics Institute, Prague (2006), hep-ph/0606300.